arXiv:0809.2864v1 [math-ph] 17 Sep 2008

New solutions for the modified generalized Degasperis–Procesi equation

Alvaro H. Salas
Department of Mathematics
Universidad de Caldas, Manizales, Colombia.
Universidad Nacional de Colombia, Manizales.
email : asalash2002@yahoo.com

José Gonzalo Escobar Lugo
Universidad Libre, Bogotá, Colombia.
Universidad Cooperativa de Colombia, Bogotá.
email : jogoel@gmail.com

Abstract. In this paper we show some new exact solutions for the generalized modified Degasperis−Procesi equation (mDP equation) $u_t - u_{xxt} + (b+1)u^2u_x = bu_xu_{xx} + uu_{xxx}$.

Keywords and phrases. Extended tanh method, nonlinear ordinary differential equation, nonlinear partial differential equation, third order evolution equation, DegasperisProcesi equation, DP equation, mDP equation, Camassa-Holm equation, CH equation, mCH equation, traveling wave solution, partial differential equation (PDE), nonlinear evolution equation.

2000 Mathematics Subject Classification. 35C05.

1. Introduction

Degasperis and Procesi [1] showed, using the method of asymptotic integrability, that the PDE

$$u_t - u_{xxt} + (b+1)uu_x = bu_xu_{xx} + uu_{xxx} \tag{1.1}$$

cannot be completely integrable unless $b = 2$ or $b = 3$. The case $b = 2$ is the Camassa Holm (CH) shallow water equation [2], which is well known to be integrable and to possess multi-soliton (weak) solutions with peaks, so called multi-peakons. Degasperis, Holm and Hone [4, 5] proved that the case $b = 3$, which they called the Degasperis Procesi (DP) equation, is also integrable and admits multi-peakon solutions. They found the two-peakon solution explicitly by direct computation.

In this paper we show some exact solutions for the generalized mDP equation [7, 8]

$$u_t - u_{xxt} + (b+1)u^2u_x = bu_xu_{xx} + uu_{xxx} \tag{1.2}$$

for any $b \neq -1$ and $b \neq -2$ by using three distinct methods: The Cole-Hopf transformation, the rational hyperbolic ansatz and the improved tanh-coth method.

2. The Cole-Hopf transformation

We seek solutions to (1.2) in the form of a generalized Cole-Hopf transformation [3][6]

$$u = A\frac{\partial^2}{\partial x^2}\log(1+\varphi(x,t)) + B, \tag{2.1}$$

for the special choice

$$\varphi(x,t) = \exp(\mu x + \lambda t + \delta), \quad (A \neq 0, \quad \lambda \neq 0 \quad \text{and} \quad \mu \neq 0.) \tag{2.2}$$

From (2.1) and (2.2) it follows that solutions have the form

$$u = \frac{A\mu^2}{2(1+\cosh(\mu\, x + \lambda\, t))} + B. \tag{2.3}$$

Applying this transformation and defining $\zeta = \exp(kx+\omega t+\delta)$, we obtain a polynomial equation in the variable ζ. Equating the coefficients of the different powers of ζ to zero, we obtain the following algebraic system :

- $B\mu^3 + \lambda\mu^2 - \left(bB^2 + B^2\right)\mu - \lambda = 0.$
- $B\mu^3 + \lambda\mu^2 + \left(bB^2 + B^2\right)\mu + \lambda = 0.$
- $(bA + A)\mu^5 - (2AB + 2AbB + 9B)\mu^3 - 9\lambda\mu^2 - \left(3bB^2 + 3B^2\right)\mu - 3\lambda = 0.$
- $(bA + A)\mu^5 + (2AB + 2AbB + 9B)\mu^3 + 9\lambda\mu^2 + \left(3bB^2 + 3B^2\right)\mu + 3\lambda = 0.$
- $\left(bA^2 + A^2 + 5bA + 11A\right)\mu^5 + (2AB + 2AbB + 10B)\mu^3 + 10\lambda\mu^2 + \left(2bB^2 + 2B^2\right)\mu + 2\lambda = 0.$
- $\left(bA^2 + A^2 + 5bA + 11A\right)\mu^5 + (2AB + 2AbB + 10B)\mu^3 + 10\lambda\mu^2 + \left(2bB^2 + 2B^2\right)\mu + 2\lambda = 0.$

Solving this system gives following solutions :

- $A = -\frac{6(b+2)}{b+1},\ B = \frac{2\mu^2-1+b\left(\mu^2-1\right)+\sqrt{1-b(b+2)\left(\mu^4-1\right)}}{2(b+1)},\ \lambda = -\frac{1}{2}\mu\left(b+1-\sqrt{1-b(b+2)\left(\mu^4-1\right)}\right)$:

$$\begin{aligned} u_1(x,t) \quad &= \quad \frac{2\mu^2-1+b\left(\mu^2-1\right)+\sqrt{1-b(b+2)\left(\mu^4-1\right)}}{2(b+1)} - \\ &\quad \frac{6(b+2)\mu^2}{2(b+1)\left(1+\cosh\left(\mu\, x - \frac{1}{2}\left(b+1-\sqrt{1-b(b+2)\left(\mu^4-1\right)}\right)\mu\, t\right)\right)}. \end{aligned}$$

- $A = -\frac{6(b+2)}{b+1},\ B = \frac{b\mu^2+2\mu^2-1-b-\sqrt{1-b(b+2)\left(\mu^4-1\right)}}{2(b+1)},\ \lambda = -\frac{1}{2}\mu\left(b+1+\sqrt{1-b(b+2)\left(\mu^4-1\right)}\right)$:

$$\begin{aligned} u_2(x,t) = &\frac{b\mu^2+2\mu^2-1-b-\sqrt{1-b(b+2)\left(\mu^4-1\right)}}{2(b+1)} - \\ &\frac{6(b+2)\mu^2}{2(b+1)\left(1+\cosh\left(\mu\, x - \frac{1}{2}\left(b+1+\sqrt{1-b(b+2)\left(\mu^4-1\right)}\right)\mu\, t\right)\right)}. \end{aligned}$$

3. The rational hyperbolic ansatz

According this method, we seek solutions in the form

$$u(x,t) = u(\xi) = \frac{a_0 + a_1\sinh(\xi) + a_2\cosh(\xi)}{1 + c_1\sinh(\xi) + c_2\cosh(\xi)}, \tag{3.1}$$

where $\xi = x + \lambda t$ and λ, a_0, a_1, a_2, c_1 and c_2 are constants to be determined. Changing the hyperbolic functions $\sinh(\xi)$ and $\cosh(\xi)$ to their exp form and substituting the resulting expression into (1.2), we obtain a polynomial equation in the variable $\zeta = \exp(\xi)$. Equating the coefficients of the powers of ζ to zero, we obtain an algebraic system in the variables λ, a_0, a_1, a_2, c_1 and c_2. After solving this system, we get the following solutions.

- $\lambda = -\frac{b}{2},\ a_0 = -\frac{3b+5}{b+1},\ a_1 = 0,\ a_2 = \frac{1}{b+1},\ c_1 = 0,\ c_2 = 1:$

$$u_3(x,t) = -\frac{3b+5-\cosh\left(x-\frac{bt}{2}\right)}{(b+1)\left(1+\cosh\left(x-\frac{bt}{2}\right)\right)}.$$

- $\lambda = -\frac{b}{2},\ a_0 = -\frac{3b+5}{b+1},\ a_1 = 0,\ a_2 = -\frac{1}{b+1},\ c_1 = 0,\ c_2 = -1:$

$$u_4(x,t) = -\frac{3b+5+\cosh\left(x-\frac{bt}{2}\right)}{(b+1)\left(1-\cosh\left(x-\frac{bt}{2}\right)\right)}.$$

- $\lambda = -\frac{b}{2}-1,\ a_0 = -\frac{3(b+2)}{b+1},\ a_1 = 0,\ a_2 = 0,\ c_1 = 0,\ c_2 = -1:$

$$u_5(x,t) = -\frac{3(b+2)}{(b+1)\left(1-\cosh\left(x-(1+\frac{b}{2})t\right)\right)}.$$

- $\lambda = -\frac{b}{2}-1,\ a_0 = -\frac{3(b+2)}{b+1},\ a_1 = 0,\ a_2 = 0,\ c_1 = 0,\ c_2 = 1:$

$$u_6(x,t) = -\frac{3(b+2)}{(b+1)\left(1+\cosh\left(x-(1+\frac{b}{2})t\right)\right)}.$$

- $\lambda = -\frac{b}{2},\ a_0 = -\frac{3b+5}{b+1},\ a_1 = -\frac{\sqrt{(b+1)^2a_2^2-1}}{b+1},\ c_2 = a_2(b+1),\ c_1 = -\sqrt{(b+1)^2a_2^2-1}:$

$$u_7(x,t) = \frac{-3b-5+(b+1)\cosh\left(x-\frac{bt}{2}\right)a_2-\sqrt{(b+1)^2a_2^2-1}\,\sinh\left(x-\frac{bt}{2}\right)}{(b+1)\left(1-\sqrt{(b+1)^2a_2^2-1}\,\sinh\left(x-\frac{bt}{2}\right)+(b+1)a_2\cosh\left(x-\frac{bt}{2}\right)\right)}.$$

- $\lambda = -\frac{b}{2},\ a_0 = -\frac{3b+5}{b+1},\ a_1 = \frac{\sqrt{(b+1)^2a_2^2-1}}{b+1},\ c_2 = a_2(b+1),\ c_1 = \sqrt{(b+1)^2a_2^2-1}:$

$$u_8(x,t) = \frac{-3b-5+(b+1)\cosh\left(x-\frac{bt}{2}\right)a_2+\sqrt{(b+1)^2a_2^2-1}\,\sinh\left(x-\frac{bt}{2}\right)}{(b+1)\left(1+\sqrt{(b+1)^2a_2^2-1}\,\sinh\left(x-\frac{bt}{2}\right)+(b+1)a_2\cosh\left(x-\frac{bt}{2}\right)\right)}.$$

- $\lambda = -\frac{b}{2}-1,\ a_0 = -\frac{3(b+2)}{b+1},\ a_1 = 0,\ a_2 = 0,\ c_1 = \sqrt{c_2^2-1}:$

$$u_9(x,t) = -\frac{3(b+2)}{(b+1)\left(1+\sqrt{c_2^2-1}\,\sinh\left(x-(1+\frac{b}{2})t\right)+c_2\cosh\left(x-(1+\frac{b}{2})t\right)\right)}.$$

- $\lambda = -\frac{b}{2}-1,\ a_0 = -\frac{3(b+2)}{b+1},\ a_1 = 0,\ a_2 = 0,\ c_1 = -\sqrt{c_2^2-1}:$

$$u_{10}(x,t) = \frac{3(b+2)}{(b+1)\left(1+\sqrt{c_2^2-1}\,\sinh\left(x-(1+\frac{b}{2})t\right)-c_2\cosh\left(x-(1+\frac{b}{2})t\right)\right)}.$$

4. The improved tanh-coth method

The traveling wave transformation

$$\xi = x + \lambda t \tag{4.1}$$

reduces (1.1) to the nonlinear ode

$$(b+1)u'(\xi)u(\xi)^2 - u'''(\xi)u(\xi) - \lambda u'''(\xi) + \lambda u'(\xi) - bu'(\xi)u''(\xi) = 0. \tag{4.2}$$

We seek solutions to (4.2) in the form

$$u(\xi) = a_0 + \sum_{i=1}^{m} \left(a_i \varphi^i + c_i \varphi^{-i}\right), \tag{4.3}$$

where m is a positive integer that will be determined and $\varphi = \varphi(\xi)$ satisfies the Riccati equation

$$\varphi'(\xi) = \alpha + \beta\varphi(\xi) + \gamma\varphi(\xi)^2, \tag{4.4}$$

where α, β, γ are constants. The solutions of (4.4) are given by:

(1) If $\alpha = 0$ and $\beta \neq 0$,

$$\varphi(\xi) = \frac{\beta}{-\gamma + \beta e^{-\beta\xi}}. \tag{4.5}$$

(2) If $\beta = \alpha = 0$ and $\gamma \neq 0$,

$$\varphi(\xi) = -\frac{1}{\gamma\xi}. \tag{4.6}$$

(3) If $\gamma = 0$ and $\beta \neq 0$,

$$\varphi(\xi) = \frac{-\alpha + \beta e^{\beta\xi}}{\beta}. \tag{4.7}$$

(4) If $\beta = 0$ and $\alpha\gamma \neq 0$,

$$\begin{cases} \varphi(\xi) = \frac{\sqrt{\alpha\gamma}}{\gamma}\tan(\sqrt{\alpha\gamma}\xi), & \alpha > 0, \gamma > 0 \\ \varphi(\xi) = \frac{\sqrt{\alpha\gamma}}{\gamma}\tanh(\sqrt{\alpha\gamma}\xi), & \alpha > 0, \gamma < 0 \\ \varphi(\xi) = \frac{\sqrt{-\alpha\gamma}}{\gamma}\tanh(-\sqrt{-\alpha\gamma}\xi), & \alpha < 0, \gamma > 0 \\ \varphi(\xi) = \frac{\sqrt{\alpha\gamma}}{\gamma}\tan(-\sqrt{\alpha\gamma}\xi), & \alpha < 0, \gamma < 0. \end{cases} \tag{4.8}$$

(5) If $\beta \neq 0$ and $\beta^2 = 4\alpha\gamma$,

$$\varphi(\xi) = -\frac{2\alpha(\beta\xi + 2)}{\beta^2\xi} \tag{4.9}$$

(6) If $\beta^2 < 4\alpha\gamma$,

$$\varphi(\xi) = \frac{\sqrt{4\alpha\gamma - \beta^2}\tan\left(\frac{1}{2}\sqrt{4\alpha\gamma - \beta^2}\xi\right) - \beta}{2\gamma}. \tag{4.10}$$

(7) If $\beta^2 > 4\alpha\gamma$ and $\gamma \neq 0$,

$$\varphi(\xi) = \frac{\sqrt{\beta^2 - 4\alpha\gamma}\tanh\left(\frac{1}{2}\sqrt{\beta^2 - 4\alpha\gamma}\xi\right) - \beta}{2\gamma}. \tag{4.11}$$

Substituting (4.3), along with (4.4) into the left hand of (4.2) and collecting all terms with the same power in $\varphi(\xi)$, we get a polynomial in the variable $\varphi = \varphi(\xi)$. This polynomial has the form

$$p\varphi^{3m+1}(\xi) + q\varphi^{2m+1}(\xi) + r\varphi^{m+3}(\xi) + \text{lower degree terms} \tag{4.12}$$

With m determined, we equate each coefficient of the polynomial to zero. This will give an overdetermined system of algebraic equations involving the parameters λ, a_0, a_i and c_i $(i = 0, \dots, m)$. Having determined these parameters, and using (4.3) and (4.5)-(4.11), we obtain analytic solutions $u(x,t)$ in a closed form.
To determine the parameter m, we usually balance the terms of highest order in the resulting polynomial. In our case, we determine m from equations $3m+1 = 2m+1$, $3m+1 = m+1$, $2m+1 = m+3$. This gives $m = 0$ or $m = 2$. We define $m = 2$ to avoid trivial solutions, so solutions have the form

$$u(\xi) = a_0 + a_1\varphi(\xi) + \frac{c_1}{\varphi(\xi)} + a_2\varphi^2(\xi) + \frac{c_2}{\varphi^2(\xi)}. \tag{4.13}$$

After solving the algebraic system , following solutions are obtained :

First Case : $\beta \neq 0$ and $\beta^2 = 4\alpha\gamma$.

- $a_0 = \frac{3(b+2)\beta^2}{2(b+1)} - 1,\ a_1 = 0,\ a_2 = 0,\ c_1 = \frac{6(b+2)\alpha\beta}{b+1},\ c_2 = \frac{6(b+2)\alpha^2}{b+1},\ \lambda = -b-1$:

$$u_{11}(x,t) = \frac{6(b+2)\beta^2}{(b+1)(\beta\xi+2)^2} - 1.$$

Second Case : $\alpha = 0$ and $\beta \neq 0$.

- $a_0 = -\frac{-b\beta^2-2\beta^2+b+\sqrt{1-b(b+2)(\beta^4-1)}+1}{2(b+1)},\ a_1 = \frac{6(b+2)\beta\gamma}{b+1},\ a_2 = \frac{6(b+2)\gamma^2}{b+1},\ c_1 = 0,\ c_2 = 0,$
$\lambda = \frac{1}{2}\left(-b - \sqrt{1-b(b+2)\left(\beta^4-1\right)} - 1\right)$:

$$\begin{aligned} u_{12}(x,\ t) \ &= \ \frac{1}{2(b+1)}\left((b+2)\beta^2 - b - 1 - \sqrt{1-b(b+2)\left(\beta^4-1\right)}\right) + \\ &\quad \frac{6(b+2)\gamma\beta^2}{b+1}\left(\frac{1}{e^{-\beta\xi}\beta-\gamma} + \frac{\gamma}{\left(e^{-\beta\xi}\beta-\gamma\right)^2}\right). \end{aligned}$$

- $a_0 = \frac{2\beta^2+b\left(\beta^2-1\right)+\sqrt{1-b(b+2)(\beta^4-1)}-1}{2(b+1)}, a_1 = \frac{6(b+2)\beta\gamma}{b+1},\ a_2 = \frac{6(b+2)\gamma^2}{b+1},\ c_1 = 0,\ c_2 = 0,$
$\lambda = \frac{1}{2}\left(-b + \sqrt{1-b(b+2)\left(\beta^4-1\right)} - 1\right)$:

$$\begin{aligned} u_{13}(x,\ t) \ &= \ \frac{1}{2(b+1)}\left((b+2)\beta^2 - b - 1 + \sqrt{1-b(b+2)\left(\beta^4-1\right)}\right) + \\ &\quad \frac{6(b+2)\gamma\beta^2}{b+1}\left(\frac{1}{e^{-\beta\xi}\beta-\gamma} + \frac{\gamma}{\left(e^{-\beta\xi}\beta-\gamma\right)^2}\right). \end{aligned}$$

Third Case : $\beta = 0$ and $\alpha\gamma \neq 0$:

- $a_0 = -\frac{-8\alpha\gamma b+b-16\alpha\gamma+\sqrt{b(b+2)(1-256\alpha^2\gamma^2)+1}+1}{2(b+1)},\ a_1 = 0\ ,\ a_2 = \frac{6(b+2)\gamma^2}{b+1},\ c_1 = 0,\ c_2 = \frac{6(b+2)\alpha^2}{b+1},$

$\lambda = \frac{1}{2}\left(-b - \sqrt{b(b+2)\left(1-256\alpha^2\gamma^2\right)+1} - 1\right)$:

$$u_{14}(x,t) = \frac{-(b+1) - 16\alpha\gamma(b+2) - \sqrt{b(b+2)\left(1-256\alpha^2\gamma^2\right)+1} + 48\alpha\gamma(b+2)\csc^2\left(2\sqrt{\alpha\gamma}\xi\right)}{2(b+1)}.$$

- $a_0 = \frac{8\alpha\gamma b - b + 16\alpha\gamma + \sqrt{b(b+2)(1-256\alpha^2\gamma^2)+1} - 1}{2(b+1)}$, $a_1 = 0$, $a_2 = \frac{6(b+2)\gamma^2}{b+1}$, $c_1 = 0$, $c_2 = \frac{6(b+2)\alpha^2}{b+1}$,
$\lambda = \frac{1}{2}\left(-b + \sqrt{b(b+2)\left(1-256\alpha^2\gamma^2\right)+1} - 1\right)$:

$$u_{15}(x,t) = \frac{-(b+1) - 16\alpha\gamma(b+2) + \sqrt{b(b+2)\left(1-256\alpha^2\gamma^2\right)+1} + 48\alpha\gamma(b+2)\csc^2\left(2\sqrt{\alpha\gamma}\xi\right)}{2(b+1)}.$$

- $a_0 = -\frac{-8\alpha\gamma b + b - 16\alpha\gamma + \sqrt{b(b+2)(1-16\alpha^2\gamma^2)+1} + 1}{2(b+1)}$, $a_1 = 0$, $a_2 = 0$, $c_1 = 0$, $c_2 = \frac{6(b+2)\alpha^2}{b+1}$,
$\lambda = \frac{1}{2}\left(-b - \sqrt{b(b+2)\left(1-16\alpha^2\gamma^2\right)+1} - 1\right)$:

$$u_{16}(x,t) = \frac{-(b+1) + 8\alpha\gamma(b+2) - \sqrt{b(b+2)\left(1-16\alpha^2\gamma^2\right)+1} + 12\alpha\gamma(b+2)\cot^2\left(\sqrt{\alpha\gamma}\xi\right)}{2(b+1)}.$$

- $a_0 = -\frac{-8\alpha\gamma b + b - 16\alpha\gamma + \sqrt{b(b+2)(1-16\alpha^2\gamma^2)+1} + 1}{2(b+1)}$, $a_1 = 0$, $a_2 = \frac{6(b+2)\gamma^2}{b+1}$, $c_1 = 0$, $c_2 = 0$,
$\lambda = \frac{1}{2}\left(-b - \sqrt{b(b+2)\left(1-16\alpha^2\gamma^2\right)+1} - 1\right)$:

$$u_{17}(x,t) = \frac{-(b+1) + 8\alpha\gamma(b+2) - \sqrt{b(b+2)\left(1-16\alpha^2\gamma^2\right)+1} + 12\alpha\gamma(b+2)\tan^2\left(\sqrt{\alpha\gamma}\xi\right)}{2(b+1)}.$$

- $a_0 = \frac{8\alpha\gamma b - b + 16\alpha\gamma + \sqrt{b(b+2)(1-16\alpha^2\gamma^2)+1} - 1}{2(b+1)}$, $a_1 = 0$, $a_2 = 0$, $c_1 = 0$, $c_2 = \frac{6(b+2)\alpha^2}{b+1}$,
$\lambda = \frac{1}{2}\left(-b + \sqrt{b(b+2)\left(1-16\alpha^2\gamma^2\right)+1} - 1\right)$:

$$u_{18}(x,t) = \frac{-(b+1) + 8\alpha\gamma(b+2) + \sqrt{b(b+2)\left(1-16\alpha^2\gamma^2\right)+1} + 12\alpha\gamma(b+2)\cot^2\left(\sqrt{\alpha\gamma}\xi\right)}{2(b+1)}.$$

- $a_0 = \frac{8\alpha\gamma b - b + 16\alpha\gamma + \sqrt{b(b+2)(1-16\alpha^2\gamma^2)+1} - 1}{2(b+1)}$, $a_1 = 0$, $a_2 = \frac{6(b+2)\gamma^2}{b+1}$, $c_1 = 0$, $c_2 = 0$,
$\lambda = \frac{1}{2}\left(-b + \sqrt{b(b+2)\left(1-16\alpha^2\gamma^2\right)+1} - 1\right)$:

$$u_{19}(x,t) = \frac{-(b+1) + 8\alpha\gamma(b+2) + \sqrt{b(b+2)\left(1-16\alpha^2\gamma^2\right)+1} + 12\alpha\gamma(b+2)\tan^2\left(\sqrt{\alpha\gamma}\xi\right)}{2(b+1)}.$$

Fourth Case : $\Delta \neq 0$, where $\Delta = \beta^2 - 4\alpha\gamma$:

- $a_0 = \frac{24\alpha\gamma + 2\Delta + b(12\alpha\gamma + \Delta - 1) - \sqrt{1 - b(b+2)(\Delta^2-1)} - 1}{2(b+1)}$, $a_1 = \frac{6(b+2)\beta\gamma}{b+1}$, $a_2 = \frac{6(b+2)\gamma^2}{b+1}$, $c_1 = 0$, $c_2 = 0$,
$\lambda = \frac{1}{2}\left(-b - \sqrt{1 - b(b+2)\left(\Delta^2-1\right)} - 1\right)$:

$$u_{20}(x,t) = -\frac{2\Delta(b+2) + b + 1 + \sqrt{1 - b(b+2)\left(\Delta^2-1\right)}}{2(b+1)} + \frac{3(b+2)\Delta}{2(b+1)}\tanh^2\left(\frac{1}{2}\sqrt{\Delta}\xi\right).$$

- $a_0 = \frac{24\alpha\gamma+2\Delta+b(12\alpha\gamma+\Delta-1)+\sqrt{1-b(b+2)(\Delta^2-1)}-1}{2(b+1)}$, $a_1 = \frac{6(b+2)\beta\gamma}{b+1}$, $a_2 = \frac{6(b+2)\gamma^2}{b+1}$, $c_1 = 0$, $c_2 = 0$, $\lambda = \frac{1}{2}\left(-b+\sqrt{1-b(b+2)\left(\Delta^2-1\right)}-1\right)$:

$$u_{21}(x,t) = -\frac{2\Delta(b+2)+b+1-\sqrt{1-b(b+2)\left(\Delta^2-1\right)}}{2(b+1)} + \frac{3(b+2)\Delta}{2(b+1)}\tanh^2\left(\frac{1}{2}\sqrt{\Delta}\xi\right).$$

- $a_0 = \frac{24\alpha\gamma+2\Delta+b(12\alpha\gamma+\Delta-1)+\sqrt{1-b(b+2)(\Delta^2-1)}-1}{2(b+1)}$, $a_1 = 0$, $a_2 = 0$, $c_1 = \frac{6(b+2)\alpha\beta}{b+1}$, $c_2 = \frac{6(b+2)\alpha^2}{b+1}$, $\lambda = \frac{1}{2}\left(-b+\sqrt{1-b(b+2)\left(\Delta^2-1\right)}-1\right)$:

$$\begin{aligned} u_{22}(x,t) &= \frac{1}{2(b+1)}((\Delta+12\alpha\gamma)(b+2)-(b+1)+\sqrt{1-b(b+2)\left(\Delta^2-1\right)}) - \\ &\quad \frac{12(b+2)\alpha\gamma\left(\beta^2+\sqrt{\Delta}\tanh\left(\frac{1}{2}\sqrt{\Delta}\xi\right)\beta-2\alpha\gamma\right)}{(b+1)\left(\beta+\sqrt{\Delta}\tanh\left(\frac{1}{2}\sqrt{\Delta}\xi\right)\right)^2}. \end{aligned}$$

- $a_0 = \frac{24\alpha\gamma+2\Delta+b(12\alpha\gamma+\Delta-1)-\sqrt{1-b(b+2)(\Delta^2-1)}-1}{2(b+1)}$, $a_1 = 0$, $a_2 = 0$, $c_1 = \frac{6(b+2)\alpha\beta}{b+1}$, $c_2 = \frac{6(b+2)\alpha^2}{b+1}$, $\lambda = \frac{1}{2}\left(-b-\sqrt{1-b(b+2)\left(\Delta^2-1\right)}-1\right)$:

$$\begin{aligned} u_{23}(x,t) &= \frac{1}{2(b+1)}\left((\Delta+12\alpha\gamma)(b+2)-(b+1)-\sqrt{1-b(b+2)(\Delta^2-1)}\right) - \\ &\quad \frac{12(b+2)\alpha\gamma\left(\beta^2+\sqrt{\Delta}\tanh\left(\frac{1}{2}\sqrt{\Delta}\xi\right)\beta-2\alpha\gamma\right)}{(b+1)\left(\beta+\sqrt{\Delta}\tanh\left(\frac{1}{2}\sqrt{\Delta}\xi\right)\right)^2}. \end{aligned}$$

5. CONCLUSIONS

In this paper, by using three distinct methods and the help of a symbolic computation engine, we obtain some exact solutions for the equation (1.1). The methods we employ certainly work well for a large class of very interesting nonlinear equations. The main advantage of these methods is that their capability of greatly reducing the size of computational work compared to existing techniques such as the pseudo spectral method, the inverse scattering method, Hirota's bilinear method, and the truncated Painlevé expansion. We think that some of the solutions we obtained are new in the literature.

REFERENCES

[1] DEGASPERIS A AND PROCESI M, *Asymptotic integrability Symmetry and Perturbation Theory*, (Rome, 1998) ed A Degasperis and G Gaeta (River Edge, NJ: World Scientific Publishing), pp. 23 - 37.

[2] CAMASSA R. AND HOLM D., *An integrable shallow water equation with peaked solitons*, Phys. Rev. Lett. **71**(1993) 16614.

[3] J. D. COLE, A quasi-linear parabolic equation occuring in aerodynamics,*Quart. Appl. Math.*, 1951, 9, pages 225-236.

[4] DEGASPERIS A., HOLM D. D. AND HONE A. N. W., *A new integrable equation with peakon solutions Theoretical and Mathematical Physics*, **133**(2002) 14631474 Preprint nlin.SI/0205023.

[5] DEGASPERIS A., HOLM D. D. AND HONE A. N. W. , *Integrable and non-integrable equations with peakons Nonlinear Physics*, Theory and Experiment (Gallipoli, 2002), vol II eds M.J. Ablowitz, M. Boiti, F. Pempinelli and B. Prinari (Singapore: World Scientific Publishing) pp 3743. Preprint nlin.SI/0209008.

[6] E. Hopf, The partial differential equation $u_t + uu_x = u_{xx}$, *Comm. Pure Appl. Math.*, **3**, 1950, 201-230.

[7] Liu Z. and Ouyang Z., *A note on solitary waves for modified forms of CamassaHolm and DegasperisProcesi equations*, Physics Letters A, **366**(2007) 377 - 381.

[8] Wazwaz A., *Solitary wave solutions for modified forms of DegasperisProcesi and CamassaHolm equations*, Physics Letters A, **352**(2006) 500 - 504.

[9] M.A. Abdou , A.A. Soliman, Modified extended tanh-function method and its application on nonlinear physical equations, Physics Letters A 353 (2006) 487 - 492.